

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Gene expression profile associated with thymus regeneration in dexamethasone-treated beef cattle**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1673655> since 2020-07-08T15:43:16Z

*Published version:*

DOI:10.1016/j.domaniend.2018.07.002

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

1 **Gene expression profile associated with thymus regeneration in dexamethasone-**  
2 **treated beef cattle**

3

4 F.T. Cannizzo, L. Starvaggi Cucuzza\*, S. Divari, E. Berio, F.E. Scaglione, B. Biolatti

5

6 *Dipartimento di Scienze Veterinarie, University of Turin, Largo Paolo Braccini 2, 10095*

7 *Grugliasco (TO), Italy*

8

9 \*Corresponding author. Tel. +39 011 670 9037;

10 Fax +39 011 6703031

11 E-mail address: laura.starvaggicucuzza@unito.it (L. Starvaggi Cucuzza)

12

## Abstract

Glucocorticoids are illegally used as growth promoters in cattle and the analytical methods officially applied most likely underestimate the precise frequency of the abuse. As a side effect, the administration of glucocorticoids causes fat infiltration, apoptosis and atrophy of the thymus. However, gross and histological observations carried out previously showed that the thymus preserves an intrinsic ability to regenerate. The aim of this work was to study the transcriptional effects of glucocorticoids on genes likely involved in regeneration of the epithelial cell network in cervical and thoracic thymus of beef cattle treated with dexamethasone or prednisolone in comparison with the control group. Moreover, the ratio of BAX/BCL2 genes was examined to verify a possible anti-apoptotic activity occurring at the same time. In cervical thymus, dexamethasone administration increased the gene expression of c-Myc ( $P < 0.01$ ), TCF3 ( $P < 0.05$ ), TP63 ( $P < 0.01$ ) and KRT5 ( $P < 0.01$ ). In thoracic thymus of dexamethasone-treated cattle the gene expression of TCF3 ( $P < 0.01$ ), TP63 ( $P < 0.01$ ) and KRT5 ( $P < 0.05$ ) was increased. These results suggested that thymic regeneration is underway in the dexamethasone-treated animals. However, the BAX/BCL2 ratio was decreased in both cervical and thoracic thymus of dexamethasone-treated cattle ( $P < 0.01$  and  $P < 0.05$ , respectively), showing an anti-apoptotic effect through the mitochondrial pathway. Conversely, prednisolone administration caused no change in the expression of all considered genes. These results sustain the hypothesis that regeneration occurs in the thymus parenchyma 6 days after the dexamethasone treatment was discontinued. This hypothesis is also supported by the absence of alterations in thymus of prednisolone-treated beef cattle. Indeed, previous studies showed the inability of prednisolone to induce macro- and microscopical lesions in the thymus. Therefore, in this context, it is not surprising that prednisolone induced no alteration of genes involved in regeneration pathway.

39    **Keywords:** beef cattle;dexamethasone; glucocorticoids; prednisolone; regeneration;  
40    thymus

## 41    **1. Introduction**

42    The administration of natural and synthetic hormones as growth promoters in animals is  
43    banned in the European Union (EU) and analytical methods are officially applied by  
44    national surveillance programs to prevent the illegal use of hormones [1].

45    Dexamethasone (DEX) is one of the most commonly administered glucocorticoids (GCs),  
46    and induces fat infiltration, increases apoptosis and causes atrophy of the thymus in cattle,  
47    as a side effect [2-4]. Conversely, prednisolone (PRD), another illicitly used GC, seems to  
48    be unable to induce thymus atrophy [5]. However, it is conceivable that the thymus  
49    preserves an intrinsic ability to regenerate after GCs administration, because the bovine  
50    thymic parenchyma and activity could be restored, as previously shown by gross and  
51    histological observations [4]. Nevertheless, the mechanisms controlling thymus  
52    regeneration remain largely unknown. Conversely, no cellular response seems to be  
53    triggered by PRD administration. It was previously shown that some transcription factors  
54    are over-expressed in the thymic stroma of mice [6]. The thymic epithelial cells (TECs)  
55    showed an up-regulation of c-Myc, TCF3 and TP63 genes during DEX- or irradiation-  
56    induced atrophy and a down-regulation after regeneration. These transcription factors  
57    were previously shown to regulate differentiation of epithelial stem cells in various tissues  
58    [7-9], suggesting a role in reconstruction/maintenance of the epithelial cell network.  
59    Moreover, it has been demonstrated that DEX- and irradiation-induced damage of the  
60    thymus resulted in proliferation of specific subset of TEC precursors expressing keratin 5  
61    (KRT5) [6].

62    Nevertheless, tissue re-growth is not only the result of cell proliferation, but also of  
63    enhanced cell survival by means of the inhibition of apoptosis or a combination of both  
64    mechanisms [10]. Several studies have highlighted that many of the molecular pathways  
65    involved in thymus atrophy rely on the mitochondria-dependent apoptotic pathway,  
66    involving proteins of the BCL2 family [11,12]. The members of BCL2 family are known to

67 be key regulatory proteins in apoptotic events, and can promote either cell survival or cell  
68 death. Indeed, the equilibrium between the pro- and anti-apoptotic members or their  
69 relative amount are crucial to sensitise the cells towards either survival or apoptosis. The  
70 anti-apoptotic effect of BCL2 acts by binding and inhibiting pro-apoptotic proteins like BAX.  
71 The latter promotes apoptosis by altering mitochondrial functions and activating the  
72 release of downstream apoptogenic factors [13]. The aim of this work was to study the  
73 biological mechanisms involved in regeneration following GCs treatment.  
74 Therefore, an increase of transcript abundance of c-Myc, TCF3, TP63, and KRT5 was  
75 hypothesized in the cervical and thoracic thymus during regeneration of the thymus in beef cattle.  
76 Additionally, the BAX and BCL2 expression and their ratio were examined to evaluate a  
77 possible anti-apoptotic activity occurring at the same time.

78

79

## 80 **2. Material and methods**

### 81 *2.1. Animals*

82 The experiment was authorized by the Italian Ministry of Health and the Ethics Committee  
83 of the University of Turin. The carcasses of the treated animals were appropriately  
84 destroyed (2003/74/CE–DL 16 March 2006, No. 158).

85 All groups of experimental animals were kept in separate pens of 10 × 15 m and were fed  
86 a diet consisting of corn silage, corn, hay and a commercial protein supplement; animals  
87 had *ad libitum* access to water. Eighteen male Charolaise beef cattle of 17 to 22 mo of age  
88 were divided into the following three groups: group A (n = 6) was administered  
89 dexamethasone–21–sodium-phosphate 0.7 mg/d *per os* for 40 d; group B (n = 6) was  
90 administered PRD 15 mg/d *per os* for 30 d; group K (n = 6) served as a control. Each  
91 morning, before the distribution of feed, the animals were tied to the feed trough, and two  
92 trained technicians administered orally one capsule containing the compound using a

drenching gun. The control animals were treated with a placebo. The animals were slaughtered 6 d after drug withdrawal. The gross and microscopic findings in the thymus of these animals were previously reported [5].

## 2.2. Samples

Thoracic and cervical thymus samples from each animal were collected and placed in RNAlater solution (Ambion, Thermo Fisher Scientific, Waltham, MA) to preserve the RNA integrity for molecular investigation.

## 2.3. Total RNA extraction and quantitative PCR

The expression of c-Myc, TCF3, TP63, KRT5, BAX and BCL2 in the thoracic and cervical thymus was investigated by quantitative PCR (qPCR). For this purpose, fifty milligrams of thymus were disrupted using a TissueLyser II (Qiagen, Hilden, Germany) with stainless steel beads in 1 mL of TRIzol (Invitrogen, Thermo Fisher Scientific). Total RNA was purified from any residual genomic DNA with a DNA-free kit (Ambion). The integrity of the RNA was confirmed by the Experion Automated Electrophoresis Station (Bio-Rad, Hercules, CA), and the concentration was measured by a spectrophotometry. cDNA was synthesised from 1 µg of total RNA using ImProm-II reverse transcriptase (Promega, Madison, WI) and random primers (Promega). To determine the amount of the specific target genes, cDNA was subjected to qPCR using the SYBRGreen method and the IQ5 (Bio-Rad) detection system. The primer sequences were designed using Primer Express v 1.5 (Applied Biosystems, Thermo Fisher Scientific) (Table 1). The peptidylprolyl isomerase A (cyclophilin A, PPIA) gene was used as a housekeeping control gene, as previously reported [14]. The expression level of each target gene was calculated using the  $2^{-\Delta Cq}$  method, where  $\Delta Cq = Cq_{\text{target gene}} - Cq_{\text{housekeeping gene}}$  [15].

## 119 2.4. Statistical analysis

120 The data were analyzed using GraphPad InStat version 3.00 (GraphPad Inc., San Diego,  
121 CA). The analysis of c-Myc, TCF3, TP63, KRT5, BAX and BCL2 gene expression and the  
122 analysis of the ratio of BAX and BCL2 expression was performed using one-way analysis  
123 of variance (ANOVA), followed by Dunnett's post hoc-test versus the control group K. If  
124 Bartlett's test suggested that the difference between the standard deviations of each group  
125 was significant, then the nonparametric Kruskal-Wallis test with Dunn's post-test versus  
126 the control group was applied. The Grubbs test was used to reveal potential outliers. A  $P$   
127 value of  $< 0.05$  was considered statistically significant. The data are shown as the mean  
128 arbitrary units ( $2^{-\Delta Cq}$ )  $\pm$  SEM.

129

130

## 131 3. Results

132 In the cervical thymus, DEX administration (group A) increased c-Myc expression (mean of  
133 mRNA arbitrary units  $\pm$  SEM:  $9.46 \times 10^{-2} \pm 1.49 \times 10^{-2}$ ) compared with the control group K  
134 ( $4.70 \times 10^{-2} \pm 3.02 \times 10^{-3}$ ) ( $P < 0.01$ ) (Fig. 1a), TCF3 expression ( $8.74 \times 10^{-3} \pm 4.54 \times 10^{-3}$ )  
135 compared with the control group K ( $8.48 \times 10^{-4} \pm 9.88 \times 10^{-5}$ ) ( $P < 0.05$ ) (Fig. 1b), TP63  
136 expression ( $8.64 \times 10^{-2} \pm 2.71 \times 10^{-2}$ ) compared with the control group K ( $1.56 \times 10^{-2} \pm$   
137  $1.85 \times 10^{-3}$ ) ( $P < 0.01$ ) (Fig. 1c) and KRT5 expression ( $2.31 \times 10^{-1} \pm 1.99 \times 10^{-2}$ ) compared  
138 with the control group K ( $3.75 \times 10^{-2} \pm 6.20 \times 10^{-3}$ ) ( $P < 0.01$ ) (Fig. 1d). Conversely, DEX  
139 administration decreased BAX expression ( $1.54 \times 10^{-3} \pm 1.17 \times 10^{-4}$ ) compared with the  
140 control group K ( $3.44 \times 10^{-3} \pm 2.38 \times 10^{-4}$ ) ( $P < 0.01$ ) (Fig. 2a). No change in BCL2  
141 expression was observed (Fig. 2b). The administration of PRD (group B) caused no  
142 change in the expression of all considered genes (Fig. 1 and 2).



143 In the thoracic thymus, DEX administration (group A) increased TCF3 expression ( $5.68 \times$   
 144  $10^{-3} \pm 1.04 \times 10^{-3}$ ) compared with the control group K ( $2.12 \times 10^{-3} \pm 2.51 \times 10^{-4}$ ) ( $P < 0.01$ )  
 145 (Fig. 1b), TP63 expression ( $6.87 \times 10^{-2} \pm 8.70 \times 10^{-3}$ ) compared with the control group K  
 146 ( $2.95 \times 10^{-2} \pm 3.62 \times 10^{-3}$ ) ( $P < 0.01$ ) (Fig. 1c) and KRT5 expression ( $1.79 \times 10^{-2} \pm 6.59 \times 10^{-$   
 147  $3$  compared with the control group K ( $5.37 \times 10^{-2} \pm 1.88 \times 10^{-3}$ ) ( $P < 0.05$ ) (Fig. 1d). No  
 148 change in c-Myc (Fig. 1a) or BCL2 expression was observed (Fig. 2b). Conversely, DEX  
 149 administration decreased BAX expression ( $3.20 \times 10^{-3} \pm 2.67 \times 10^{-4}$ ) compared with the  
 150 control group K ( $4.91 \times 10^{-3} \pm 5.35 \times 10^{-4}$ ) ( $P < 0.05$ ) (Fig. 2a). The administration of PRD  
 151 (group B) caused no change in the expression of all considered genes (Fig. 1 and 2).  
 152 The BAX/BCL2 ratio was statistically different in both cervical and thoracic thymus of  
 153 Group A compared to the controls ( $P < 0.01$  and  $P < 0.05$ , respectively) (Fig. 2c).

154

155

#### 156 **4. Discussion**

157 Council Directive 96/22/EC [1], as amended by Directives 2003/74/EC [16] and  
 158 2008/97/EC [17], stipulates that all use of steroids,  $\beta$ -agonists or other substances for the  
 159 chemical manipulation of animal growth is severely banned in the EU. However, results  
 160 from studies conducted by the Italian Health Ministry indicate that these substances are  
 161 persistently used, and therefore a permanent commitment by the public veterinary services  
 162 for their prevention and control is required. The thymus represents a GCs target tissue,  
 163 and in vitro or ex vivo qualitative and semi-quantitative morphological investigations to  
 164 identify the cellular effects of GCs were reported [2,4,18].

165 Indeed, the thymus weight and volume of beef cattle following long-term administration of  
 166 low doses of DEX were significantly reduced compared with the controls [4]. Moreover,  
 167 DEX-treated animals showed severe thymus atrophy, characterized by a serious volume

168 reduction of the organ, which in some animals almost disappeared and was replaced by  
169 fat tissue. Histologically, the thymic cortex undergoes atrophy, while the medullary  
170 framework was still present though reduced, showing a pronounced rarefaction of  
171 lymphocytes [5]. Conversely, no histological change was observed in the thymus following  
172 long-term treatment with PRD [5].

173 However, partial recovery of thymus weight and structure after 25 days followed by  
174 complete recovery after 32 days was observed in veal calves [3] and similar results have  
175 been observed in thymus of beef cattle examined 26 days after the end of treatment [4].

176 Thus, it is conceivable that the thymus preserves an intrinsic ability to regenerate, but the  
177 molecular mechanisms controlling the regeneration of the thymus are largely unknown.

178 Previous work in mice demonstrated that c-Myc, TP63, and TCF3 gene expression was  
179 up-regulated in TECs during peak thymic atrophy and was down-regulated at later time  
180 points when thymuses were undergoing regeneration [6]. These transcription factors were  
181 previously shown to regulate differentiation of epithelial stem cells in various tissues  
182 [7,8,19] suggesting a role in reconstruction/maintenance of the epithelial cell network.

183 Consistent with these findings, our results showed an up-regulation of the same  
184 transcription factors in thymus of beef cattle experimentally treated with a low doses of  
185 DEX for a long-term. In contrast, the treatment with PRD did not induce any expression  
186 changes in the genes examined in this study. Since the DEX administration induces the  
187 thymus atrophy [4], whereas PRD treatment appears to have no effects on the thymus  
188 tissue [5], the expression of the transcription factors might play a role in regeneration of  
189 the thymic stroma.

190 Moreover, it has been demonstrated that DEX-induced damage of the thymus resulted in  
191 proliferation of subset of TEC precursors expressing KRT5 [6]. This active expansion could  
192 explain the significantly over-expression of KRT5 observed in the atrophic thymus of DEX-  
193 treated beef cattle compared to controls.

194 Glucocorticoids influence the growth and differentiation of thymocytes through several  
195 mechanisms including apoptosis [20]. The apoptotic effect of GCs could shift the balance  
196 between expression of pro-survival and pro-apoptotic factors, ultimately leading to cell  
197 death or apoptosis [21]. In this study, DEX administration caused a decrease of BAX  
198 expression, whereas BCL2 expression remain unchanged. Moreover, the ratio of BAX and  
199 BCL2 expression diminished. These results suggest that an anti-apoptotic effect is  
200 occurring through the mitochondrial pathway, and this may support the hypothesis that  
201 regeneration activity occurs in the thymus parenchyma 6 days after the treatment is  
202 discontinued.

203 Thus, our data suggested that the observed recovery of thymus of DEX-treated beef cattle  
204 might be mediated by several events, including the elevated expression of transcription  
205 factors involved in differentiation of epithelial stem cells, active proliferation of TECs subset  
206 and inhibition of apoptosis. The expression of this panel of genes appears characteristic of  
207 the animals treated with DEX and not of PRD-treated animals. Indeed, previous studies  
208 showed the inability of PRD to induce macro- and microscopical lesions (i.e. atrophy) in  
209 thymus. Therefore, it is not surprising that PRD does not induce alteration of genes  
210 involved in regeneration pathway.

211

212

## 213 **Acknowledgments**

214 This work was partially funded by the Ministero delle Politiche Agricole Alimentari e  
215 Forestali SAFORISK project “Prevenzione dell’uso di anabolizzanti in zootecnia.  
216 Creazione di un Marchio a difesa degli allevamenti italiani” (D.M. 2089/09, 29th of January  
217 2009). The authors are grateful to Dr. Sara Baldizzone.

218 Declarations of interest: none.

219

220

## 221 **References**

222 [1] Council Directive 96/23/EC of 29 April 1996 on measures to monitor certain substances  
223 and residues thereof in live animals and animal products and repealing Directives  
224 85/358/EEC and 86/469/EEC and Decisions 89/187/EEC and 91/664/EEC. Off J Eur  
225 Union 1996, L 125, 10-32.

226

227 [2] Biolatti B, Bollo E, Cannizzo FT, Zancanaro G, Tarantola M, Dacasto M, Cantiello M,  
228 Carletti M, Biolatti PG, Barbarino G. Effects of low-dose dexamethasone on thymus  
229 morphology and immunological parameters in veal calves. J Vet Med A Physiol Pathol Clin  
230 Med 2005;52:202-208.

231

232 [3] Cannizzo FT, Miniscalco B, Riondato F, Bollo E, Barbarino G, Giorgi P, Mazzini C,  
233 Biolatti B. Effects of anabolic and therapeutic doses of dexamethasone on thymus  
234 morphology and apoptosis in veal calves. Vet Rec 2008;163:448-452.

235

236 [4] Cannizzo, FT, Spada, F, Benevelli, R, Nebbia, C, Giorgi, P, Brina, N, Bollo, E, Biolatti,  
237 B, Thymus atrophy and regeneration following dexamethasone administration to beef  
238 cattle. Vet Rec 2010;167:338-343.

239

240 [5] Cannizzo FT, Capra P, Divari S, Ciccotelli V, Biolatti B, Vincenti M. Effects of low-dose  
241 dexamethasone and prednisolone long term administration in beef calf: Chemical and  
242 morphological investigation. Anal Chim Acta 2011;700:95-104.

243

244 [6] Popa I, Zubkova I, Medvedovic M, Romantseva T, Mostowski H, Boyd R, Zaitseva M.  
245 Regeneration of the adult thymus is preceded by the expansion of K5<sup>+</sup>K8<sup>+</sup> epithelial cell

246 progenitors and by increased expression of Trp63, cMyc and Tcf3 transcription factors in  
247 the thymic stroma. *Int Immunol* 2007;19:1249-1260.

248

249 [7] Merrill BJ, Gat U, DasGupta R, Fuchs E. Tcf3 and Lef1 regulate lineage differentiation of  
250 multipotent stem cells in skin. *Genes Dev* 2001;15:1688-1705.

251

252 [8] Nair M, Teng, A, Bilanchone V, Agrawal A, Li B, Dai X. Ovol1 regulates the growth arrest  
253 of embryonic epidermal progenitor cells and represses c-myc transcription. *J Cell Biol*  
254 2006;173:253-264.

255

256 [9] Yang A, Kaghad M, Wang Y, Gillett E, Fleming MD, Dötsch V, Andrews NC, Caput D  
257 McKeon F. p63, a p53 homolog at 3q27-29, encodes multiple products with  
258 transactivating, death-inducing, and dominant-negative activities. *Mol Cell* 1998;2:305-316.

259

260 [10] Hannun YA. Apoptosis and the dilemma of cancer chemotherapy. *Blood*  
261 1997;89:1845-1853

262

263 [11] Kanavaros P, Stefanaki K, Rontogianni D, Papalazarou D, Sgantzos M, Arvanitis D,  
264 Vamvouka C, Gorgoulis V, Siatitsas I, Agnantis NJ, Bai M. Immunohistochemical expression  
265 of p53, p21/waf1, rb, p16, cyclin D1, p27, Ki67, cyclin A, cyclin B1, bcl2, bax and bak proteins  
266 and apoptotic index in normal thymus. *Histol Histopathol* 2001;16:1005-1012.

267

268 [12] Korsmeyer SJ. Bcl-2: an antidote to programmed cell death. *Cancer Surv* 1992;15:105-  
269 118.

270

271 [13] Wei MC, Zong WX, Cheng EH, Lindsten T, Panoutsakopoulou V, Ross AJ, Roth KA,  
 272 MacGregor GR, Thompson CB, Korsmeyer SJ. Proapoptotic BAX and BAK: a requisite  
 273 gateway to mitochondrial dysfunction and death. *Science* 2001;292:727-730.

274

275 [14] Starvaggi Cucuzza L, Biolatti B, Scaglione FE, Cannizzo FT. Role of FKBP51 in the  
 276 modulation of the expression of the corticosteroid receptors in bovine thymus following  
 277 glucocorticoid administration. *Domest Anim Endocrinol* 2018;62:10-15.

278

279 [15] Schmittgen TD, Livak LKJ. Analyzing real-time PCR data by the comparative C<sub>T</sub>  
 280 method. *Nat Protoc* 2008;3:1101-1118.

281

282 [16] Directive 2003/74/EC of the European Parliament and of the Council of 22 September  
 283 2003 amending Council Directive 96/22/EC concerning the prohibition on the use in  
 284 stockfarming of certain substances having a hormonal or thyrostatic action and of beta-  
 285 agonists. *Off J Eur Union* 2003, L 262, 17–21.

286

287 [17] Directive 2008/97/EC of the European Parliament and of the Council of 19 November 2008  
 288 amending Council Directive 96/22/EC concerning the prohibition on the use in stockfarming of certain  
 289 substances having a hormonal or thyrostatic action and of beta-agonists. *Off J Eur Union* L 318, 9–  
 290 11.

291

292 [18] Jacysyn JF, Conde-Moscatelli M, Barrichello CR, Silva UR, Macero MS, Amarante-  
 293 Mendes GP. Thymic epithelial cells mediate a Bcl-2-independent protection of single-  
 294 positive thymocytes from dexamethasone-induced apoptosis. *Exp Cell Res* 2002;272:119-  
 295 126.

296

297 [19] Murphy MJ, Wilson A, Trumpp A. More than just proliferation: Myc function in stem cells.  
298 Trends Cell Biol 2005;15:128-137.  
299  
300 [20] Bruscoli S, Di Virgilio R, Donato V, Velardi E, Baldoni M, Marchetti C, Migliorati G,  
301 Riccardi C. Genomic and non-genomic effects of different glucocorticoids on mouse  
302 thymocyte apoptosis. Eur J Pharmacol 2006;529:63-70.  
303  
304 [21] Wang D, Muller N, Kirsty G, McPetherson KG, Reichardt HM. Glucocorticoids engage  
305 different signal transduction pathways to induce apoptosis in thymocytes and mature T  
306 cells. J Immunol 2006;176:1695-1702.

307 **Table 1** Primer sequences used for qPCR.

Gene	Sense	Sequences 5' → 3'	Amplicon size (bp)	GeneBank accession number
c-Myc	Forward	ATGCCACGTGTCTACCCATCA	100	NM_001046074
	Reverse	GACCCTGCCACTGTCCAACT		
TCF3	Forward	TGGCTGAGTGCACCCTGAA	107	XM_002691408
	Reverse	CCGGGCCAATTCGTAGTACTT		
TP63	Forward	TTCCGTGAGCCAGCTTATCA	100	NM_001191337
	Reverse	GTGGGTGCCCATCATAGGAAT		
KRT5	Forward	GAGCCTTTGTTAGAGCAGTACATCAA	100	NM_001008663
	Reverse	CCTGCATATTCCTGAGCTCTGA		
BAX	Forward	AAGCGCATCGGAGATGAATT	100	NM_173894
	Reverse	CCGCCACTCGGAAAAAGAC		
BCL2	Forward	TGGTGGGCGCTTGCAT	100	NM_001166486
	Reverse	TTCTGCTGCTTCTTGAATCTTCTG		
PPIA	Forward	GCCCCAACACAAATGGTT	95	NM_178320
	Reverse	CCCTCTTTCACCTTGCCAAAG		

308



309 **Figure captions**

310 **Fig. 1.** Effects of dexamethasone-21-sodium phosphate (DEX; group A) or prednisolone  
311 (PRD; group B) on c-Myc (a), TCF3 (b), TP63 (c) and KRT5 (d) gene expression  
312 compared with the control group K in the cervical and thoracic thymus of beef cattle. The  
313 results are presented as the means  $\pm$  SEM. The y-axes show arbitrary units representing  
314 relative mRNA expression levels. \* $P < 0.05$ , \*\* $P < 0.01$  versus the control group K.

315

316 **Fig. 2.** Effects of dexamethasone-21-sodium phosphate (DEX; group A) or prednisolone  
317 (PRD; group B) on BAX (a) and BCL2 (b) gene expression compared with the control group  
318 K in the cervical and thoracic thymus of beef cattle. The results are presented as the means  
319  $\pm$  SEM. The y-axes show arbitrary units representing relative mRNA expression levels.  
320 Effects of dexamethasone-21-sodium phosphate (DEX; group A) or prednisolone (PRD;  
321 group B) on ratio of BAX and BCL2 expression (c) in cervical and thoracic thymus of beef  
322 cattle. \* $P < 0.05$ , \*\* $P < 0.01$  versus the control group K.